

REFERENCES

1. Berry JD, Willis B, Gupta S, et al. Lifetime risks for cardiovascular disease mortality by cardiorespiratory fitness levels measured at ages 45, 55, and 65 years in men. The Cooper Center Longitudinal Study. *J Am Coll Cardiol* 2011;57:1604–10.
2. Gupta S, Rohatgi A, Ayers CR, et al. Cardiorespiratory fitness and classification of risk of cardiovascular disease mortality. *Circulation* 2011;123:1377–83.

Sex-Specific Outcomes for HeartMate II

We read with great interest the recent paper by Starling et al. (1), which provides results on the post-marketing study required by the Food and Drug Administration (FDA) as a condition of approval of the HeartMate II (Thoratec Corporation, Pleasanton, California). However, the value of this report would be strengthened considerably if the authors would provide sex-specific data for the 38 women (22% of the patient population) in the registry.

Although the FDA requires sex-specific data for all high-risk device approvals, a recent analysis found that only 41% of recent cardiovascular device Summaries of Safety and Effectiveness Data reported such data (2). The HeartMate II was approved based on data in just 44 women (3). Furthermore, the FDA's Summary of Safety and Effectiveness Data noted that women had an 18% risk of stroke versus 6% in men and trends toward greater bleeding and infection, although it was not possible to make conclusions on differences in safety between men and women (4). Therefore, the FDA specifically stated that the required post-approval study (INTERMACS [Interagency Registry for Mechanically Assisted Circulatory Support]) would collect data on "gender-specific outcomes." Given that sex-specific outcomes was an FDA condition and that the device has been specifically advertised as being suitable for women (5), we believe the authors should report sex-specific results, which would be particularly helpful for the outcomes in Figures 2 and 3 and adverse events in Table 3. Adequate sex-specific data will help to clarify any differences in outcomes between men and women (6) and enable better care of all patients with advanced heart failure.

Sanket S. Dhruva, MD

***Rita F. Redberg, MD, MSC**

*Women's Cardiovascular Services
University of California, San Francisco
505 Parnassus Avenue, Suite M-1180
San Francisco, California 94143-0124
E-mail: redberg@medicine.ucsf.edu

doi:10.1016/j.jacc.2011.05.039

REFERENCES

1. Starling RC, Naka Y, Boyle AJ, et al. Results of the post-U.S. Food and Drug Administration-approval study with a continuous flow left ventricular assist device as a bridge to heart transplantation: a prospective

study using the INTERMACS (Interagency Registry for Mechanically Assisted Circulatory Support). *J Am Coll Cardiol* 2011;57:1890–8.

2. Dhruva SS, Bero LA, Redberg RF. Gender bias in studies for Food and Drug Administration premarket approval of cardiovascular devices. *Circ Cardiovasc Qual Outcomes* 2011;4:165–71.
3. Dhruva SS, Redberg RF. The need for sex-specific data prior to Food and Drug Administration approval (letter). *J Am Coll Cardiol* 2010; 55:261.
4. Summary of Safety and Effectiveness Data. PMA 060040. Thoratec HeartMate II Left Ventricular Assist System (LVAS). April 2008. Available at: http://www.accessdata.fda.gov/cdrh_docs/pdf6/P060040b.pdf. Accessed May 6, 2011.
5. Thoratec Corporation. HeartMate II Left Ventricular Assist System: The New Era Begins. Available at: <http://www.thoratec.com/medical-professionals/vad-product-information/heartmate-ii-lvas.aspx>. Accessed May 6, 2011.
6. Hayes SN, Redberg RF. Dispelling the myths: calling for sex-specific reporting of trial results. *Mayo Clin Proc* 2008;83:523–5.

Reply

On behalf of the authors, we thank Drs. Dhruva and Redberg for their letter in reference to our paper (1). We strongly acknowledge the importance that women are well represented in clinical trials and that sex-specific outcomes are reported. Indeed, the newer axial flow pumps have now enabled a wider dissemination of left ventricular assist devices (LVADs), because prior devices were limited to patients with body surface areas $>1.5 \text{ m}^2$. We reported a consecutive group of patients implanted with the HeartMate II (HMII) (Thoratec Corporation, Pleasanton, California) device and the previously-approved Federal Drug Administration devices as per a stipulated post-market approval study.

Of the 169 HMII patients, there were 131 men (78%) and 48 women (22%). The percentage of patients who were transplanted, recovered, or received ongoing support at 180 days was 90% for men and 92% for women. A smaller percentage of women (24%) received transplants by 12 months of support compared with men (39%), and a larger percentage were still on LVAD support (63% women vs. 47% males) at 12 months. However, there was no difference in overall survival (log rank $p = 0.4038$), and the 1-year survival estimate for patients remaining on LVAD support was $83.8 \pm 6.7\%$ (women) versus $88.3 \pm 3.0\%$ (men). There were no statistically significant differences in any adverse event between women and men. The control group had a smaller percentage of women (17% vs. 22%), but this was not statistically significant ($p = 0.217$). In comparison with results in the clinical trial for bridge to transplantation for the HeartMate II LVAD in 465 patients as reported by Bogaev et al. (2), the distribution of women (22%) and 361 men (78%) was the same. In addition, similar to the post-approval study, there was a smaller percentage of women receiving heart transplants over the first 18 months of support and a greater percentage with ongoing support compared with men, and also with no difference in overall survival. In contrast to the post-approval study, in the clinical trial, hemorrhagic stroke occurred more frequently in women versus men, and device-related infections occurred less frequently than in men.

Finally, it should be noted that historically, the percentage of women undergoing cardiac transplantation has been significantly lower than men. The latest international heart transplant registry report shows: men, 80.1% from 1992 to 2001 and 77.1% from 2002 to 2006/2009; $p < 0.0001$ (3). We fully agree with Drs. Dhruva and Redberg of the importance to collect and report